

48세 여자에서 발견된 비인강 유피낭종 1예

울산대학교 의과대학 서울아산병원 이비인후과학교실,¹ 가천의과대학교 중앙길병원 이비인후과학교실²
정유삼¹ · 이정훈² · 전형규² · 최진호²

A Case of Nasopharyngeal Dermoid in an 48 Year Old Women

Yoo-Sam Chung, MD¹, Jung Hoon Lee, MD², Hyung Gyu Chun, MD² and Jin Ho Choi, MD²

¹Department of Otolaryngology, Asan Medical Center, University of Ulsan College of Medicine, Seoul,

²Department of Otolaryngology-Head and Neck Surgery, Gil Medical Center,
Gachon Medical College, Incheon, Korea

-ABSTRACT -

Dermoids are rare lesions affecting the nasopharynx. Dermoids are composed of two germinal layers, ectoderm and mesoderm. They usually diagnosed at birth or shortly thereafter. The male-to-female ratio is 1 : 6 in cases of dermoids. Symptoms are dependent on the size and location of the lesion. The patients with large lesion present dyspnea, severe difficulties in sucking, feeding immediately after birth. They are treated by simple surgical excision. Recurrences are rare. We report an unusual case of dermoid in nasopharynx appeared in a 48 year old women presenting snoring. The dermoid was treated with simple surgical excision using the intraoral approach. (J Clinical Otolaryngol 2004;15:143-145)

KEY WORDS : Nasopharynx · Dermoid cyst.

서론		1 : 6		4)	
1)	1.6~6.9%	2)	120	3)	가
1989		48		5)	
3)	1977	1			
1998	17 가	4)			
가					
증례					
: 2004	4	14			
: 2004	5	17	48		
:	138 - 736	388 - 1			164 cm
: (02) 2224 - 3710		: (02) 489 - 2773	65 kg		
E - mail : yschung@amc.seoul.kr					




Fig. 1. Enhanced nasopharyngeal computed tomogram. A : axial view shows 4x3 cm sized left side nasopharyngeal mass with central necrosis. B : coronal view.



Fig. 2. Gross finding of specimen. It had smooth surface and was soft.




Fig. 3. Microscopic finding of the mass. It shows keratinized squamous epithelium hair follicles and sebaceous gland. It was diagnosed with dermoid (H & E, x40).

가
(laser assisted uvulopalatoplasty)

4x3 cm 가

4x3x3 cm

(Fig. 1).

(Fig. 2).

(Fig. 3).

1 (glioma),
10 (congenital rhabdomyosarcoma),
(neurofibroma) ⁴⁾

고 찰

(Thornwalt’s cyst),
(extramedullary plasmacytoma)
0.5~6 cm

가 1 : 6 ⁴⁾ 20

가 73 가

가 ⁵⁾ Arnold’s

가 가 ⁶⁾ 가 ³⁾ 가

Epignathi ⁴⁾

가 48

(teratoid tumor)

가

(teratoma) Epignathi

가 가

가 가

가 가

가

가 (inclusion theory)

가 ³⁾

(totipotent)

, blastomere

stage ⁷⁾

가 가

가

가

가 ³⁾

중심 단어 :

REFERENCES

- 1) Chakravarti A, Vishwakarma SK, Arora VK, Singh I. *Dermoid (hairy polyp) of the nasopharynx. Indian J Pediatr 1998;65:473-6.*
- 2) New GB, Erich JB. *Dermoid cysts of the head and neck. Surg Gynecol Obstet 1937;65:48.*
- 3) McShane D, el Sherif I, Dotle-Kelly W, Fennel G, Walsh M. *Dermoids (‘hairy polyps’) of the oro-nasopharynx. J Laryngol Otol 1989;103:612-5.*
- 4) Coppit GL 3rd, Perkins JA, Manning SC. *Nasopharyngeal teratomas and dermoids: a review of the literature and case series. Int J Pediatr Otorhinolaryngol 2000;52:219-27.*
- 5) Resta L, Santangelo A, Lastilla G. *The s.c. ‘hairy polyp’ or ‘dermoid’ of the nasopharynx (An unusual observation in older age). J Laryngol Otol 1984;98:1043-6.*
- 6) Arnold J. *Ein Fall von congenitalem zusammengestztem Lipon der Zunge und der Pharynx mit perforation in die Schadelhoehle. Virchows Arch Pathol Anat Physiol Klin Med 1870;50:482-516.*
- 7) Holt GR, Holt JE, Weaver RG. *Dermoids and teratomas of the head and neck. Ear Nose Throat J 1979;58:520-31.*